



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## The ability of the toe-brachial index to predict the outcome of treadmill exercise testing in patients with a normal resting ankle-brachial index

Høyer, Christian; Strandberg, Jesper; Overvad Jordansen, Malene Kragh; Zacho, Helle D

*Published in:*  
Annals of Vascular Surgery

*DOI (link to publication from Publisher):*  
[10.1016/j.avsg.2019.10.041](https://doi.org/10.1016/j.avsg.2019.10.041)

*Creative Commons License*  
CC BY-NC-ND 4.0

*Publication date:*  
2020

*Document Version*  
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Høyer, C., Strandberg, J., Overvad Jordansen, M. K., & Zacho, H. D. (2020). The ability of the toe-brachial index to predict the outcome of treadmill exercise testing in patients with a normal resting ankle-brachial index. *Annals of Vascular Surgery*, 64, 263-269. <https://doi.org/10.1016/j.avsg.2019.10.041>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

# Journal Pre-proof

The ability of the toe-brachial index to predict the outcome of treadmill exercise testing in patients with a normal resting ankle-brachial index.

Christian Høyer, MD, PhD, Jesper Strandberg, MD, Malene Kragh Overvad Jordansen, BS, Helle D. Zacho, MD, PhD



PII: S0890-5096(19)30887-8

DOI: <https://doi.org/10.1016/j.avsg.2019.10.041>

Reference: AVSG 4703

To appear in: *Annals of Vascular Surgery*

Received Date: 24 June 2019

Revised Date: 24 September 2019

Accepted Date: 1 October 2019

Please cite this article as: Høyer C, Strandberg J, Overvad Jordansen MK, Zacho HD, The ability of the toe-brachial index to predict the outcome of treadmill exercise testing in patients with a normal resting ankle-brachial index., *Annals of Vascular Surgery* (2019), doi: <https://doi.org/10.1016/j.avsg.2019.10.041>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 Elsevier Inc. All rights reserved.

# The ability of the toe-brachial index to predict the outcome of treadmill exercise testing in patients with a normal resting ankle-brachial index.

Christian Høyer, MD, PhD<sup>1,2</sup>  
Jesper Strandberg, MD<sup>2</sup>  
Malene Kragh Overvad Jordansen, BS<sup>2</sup>  
Helle D Zacho, MD, PhD<sup>2</sup>

(1) Department of Clinical Physiology, Viborg Regional Hospital, DK-8800 Viborg, Denmark.  
(2) Department of Nuclear Medicine, Aalborg University Hospital, DK-9000 Aalborg, Denmark.

*Corresponding author:*  
Christian Høyer, MD, PhD, As. Prof.  
Department of Clinical Physiology  
Viborg Regional Hospital  
Heibergs Allé 4  
DK-8800 Viborg  
Denmark

Tel (+45) 7844 3300  
Fax (+45) 7844 3399  
E-mail: christian.hoyer@midt.rm.dk

E-mail for co-authors: JS: [j.strandberg@rn.dk](mailto:j.strandberg@rn.dk)  
MKOJ: [mako@rn.dk](mailto:mako@rn.dk)  
HDZ: [h.zacho@rn.dk](mailto:h.zacho@rn.dk)

Manuscript category: Original article

Word count: 2,737

Mesh terms: Ankle brachial index  
Toe brachial index  
Diagnostic test  
Peripheral arterial disease  
Postexercise ABI

## 50 **Abstract**

### 51 **Objective**

53 Peripheral arterial disease (PAD) in the presence of a normal ankle-brachial index (ABI) can be  
54 diagnosed noninvasively by measuring a postexercise ABI or by measuring the toe-brachial index  
55 (TBI).

### 56 **Methods**

57 This was a prospective comparative study. Over a period of 30 months, a total of 415 patients who  
58 were referred with the suspicion of vascular claudication and resting values of  $0.91 \leq \text{ABI} < 1.40$  were  
59 further evaluated for the resting TBI and postexercise ABI by treadmill testing.

### 60 **Results**

61 A total of 325 (39%) of the 830 investigated limbs had a low TBI ( $\leq 0.70$ ), and 505 (61%) had a  
62 normal TBI. Of the limbs with a low TBI, 160 (49%) had PAD according to a postexercise ABI  
63 versus 165 (33%) of the limbs with normal TBI. The overall agreement in PAD classification  
64 between the two methods was 500/830 (60%) with a Cohen's kappa = 0.166 (95% CI: 0.096-0.232).  
65 The data showed an inverse correlation between the magnitude of the TBI decrease, as well as the  
66 resting ABI, and the probability of an abnormal postexercise ABI. On average, limbs with a low  
67 TBI had a lower resting ABI than patients with a normal TBI ( $1.07 \pm 0.09$  vs.  $1.13 \pm 0.10$ ,  $P < .001$ ).  
68 The groups with a low TBI had a significantly higher ratio of abnormal test results than patients  
69 with a normal TBI, in limbs with ABI (0.96-1.00) and ABI  $> 1.10$  ( $P \leq .022$  for both), but there  
70 were no statistically significant differences found in other ABI intervals ( $P > .200$  for all).

### 71 **Conclusions**

72 The magnitude of the TBI reduction correlates with an increased probability of an abnormal  
73 postexercise ABI. However, this is due in part to limbs with a low TBI having a lower resting ABI  
74 on average than limbs with a normal TBI, which also correlates with the probability of an abnormal

exercise test result. This study shows that the TBI and the postexercise ABI are not interchangeable for establishing a PAD diagnosis.

Keywords: Ankle brachial index (MeSH), Toe brachial index (MeSH), Diagnostic test (MeSH), Peripheral arterial disease (MeSH), Postexercise ABI.

## 1. Introduction

Arteriosclerosis in the lower limbs is a common condition affecting 15-20% of persons older than 70-years.<sup>1</sup> The presence and severity of peripheral arterial disease (PAD) can be diagnosed by measuring the ankle-brachial index (ABI) at rest. Having an  $ABI \leq 0.90$  is indicative of the presence of arterial disease and is associated with an increased risk of cardiovascular mortality and morbidity. However, a normal resting ABI does not rule out the presence of PAD.<sup>2</sup>

The normal physiological response to leg exercise, such as walking, is an increase in the central systolic pressure along with vasodilation in the exercising muscles, thus reducing peripheral vascular resistance. In cases with occlusive PAD, this can lead to a significant drop in ankle pressure or ABI post exercise.<sup>3</sup> The resting ABI can be unreliable in conditions related to medial arterial calcinosis such as diabetes, chronic kidney insufficiency, or age, which can lead to falsely elevated, or even falsely normal ankle pressures.<sup>4</sup>

In addition to the ABI, PAD can be diagnosed by measuring the toe-brachial index (TBI), which is less prone to medial arterial calcinosis.<sup>5</sup> Previous studies have shown that approximately 20% of patients suspected of PAD have normal ankle pressures but low TBI.<sup>6</sup> However, this subset of patients consists of both patients with masked large vessel disease, and small vessel disease with lesions below the ankle level. The aim of the current study is to test the ability of the toe-brachial index to predict the outcome of treadmill testing, and establish whether a normal TBI would render a supplementary treadmill testing redundant.

## 141 2. Materials and Methods

### 142 2.1 Design

143 This was a prospective study.

### 144 2.2 Subjects

145 Over a 30 month inclusion period from 2015-2018, patients who were referred to the Department of  
146 Nuclear Medicine at Aalborg University Hospital for distal blood pressure measurements and  
147 clinically suspected of vascular claudication were screened for enrollment. The inclusion criteria  
148 were patients with  $0.91 \leq \text{ABI} < 1.40$  who were able to participate in standardized treadmill testing.  
149 The study protocol was approved by the Northern Danish Regional Committee on Biomedical  
150 Research Ethics and the Danish Data Protection Agency. The study complied with the Helsinki II  
151 declaration.

### 152 2.3 Experimental procedure

153 The patients rested in a supine position for at least 15 minutes prior to the measurements. Adequate  
154 limb temperatures were maintained at 35-40°C using heating overlays prior to testing (Dorcas  
155 Activator Pack, ProTerapi, Ballerup, Denmark). Toe- and ankle pressures were assessed by  
156 photoplethysmography (Falcon PRO, Viasonix, Ra'anana, Israel).<sup>7</sup> Appropriately sized pneumatic  
157 occlusion cuffs were positioned at the site of measurement. The photoplethysmographic sensors were  
158 positioned distal to the cuff and measured changes in light absorption upon illumination of the skin.  
159 The cuff deflated automatically with the sensor detecting flow throughout the deflation period. The  
160 brachial blood pressure was measured simultaneously with each toe or ankle pressure (automatic  
161 digital blood pressure monitor M6, Omron Healthcare Europe, Hoofddorp, Netherlands), with the  
162 side with the highest systolic pressure selected as the reference for the ABI and TBI calculations.

## 2.4 Exercise test

Following the assessment of the resting ABI and TBI, the patients underwent standardized treadmill testing at a set speed of 4 kilometers per hour (2.5 miles per hour), and at a fixed inclination (10%) for 5 minutes or until they were unable to continue due to symptoms (MedTrack CR60, Quinton Instrument, Washington, USA). Immediately following completion of the treadmill exercise, the patients were repositioned in a supine position, and ankle and brachial measurements were repeated until the normalization of the ABI. The observers were not blinded to the outcome of the TBI measurements.

## 2.5 Diagnostic classification

A resting ABI was considered within the normal range between  $0.91 \leq \text{ABI} < 1.40$ , and  $\text{TBI} > 0.70$  according to criteria set by the American College of Cardiology/American Heart Association (ACC/AHA).<sup>1,2</sup> An abnormal postexercise ABI was defined according to the AHA scientific statement by one of the two following criteria: a postexercise ABI decrease of more than 20% or a postexercise ankle pressure decrease of more than 30 mmHg.<sup>3</sup> An abnormal postexercise increase in ABI was defined as an ABI increase in either of the two limbs.<sup>8</sup> This was, however, not deemed an abnormal test result.

## 2.6 Statistical analysis

The data are presented as the mean  $\pm$  standard deviation. The differences in hemodynamic and demographic variables between the two groups were analyzed using an unpaired t-test, in the case of quantitative variables and a chi-square test ( $\chi^2$ ) in the case of categorical variables. Agreement in diagnostic classification (PAD/not PAD) was analyzed by Cohen's kappa ( $\kappa$ ). A P-value  $< 0.05$  was considered to be statistically significant, and the statistical analysis was performed using SPSS software version 20.0 (SPSS Inc., Illinois, USA).



### 3 Results

#### 3.1 Patients and data sampling

During the 30 month inclusion period, a total of 4020 distal pressure measurements were performed at the department, of which 432 patients met the inclusion criteria. A total of 18 patients were excluded due to incomplete exercise tests ( $n=15$ ) or failure to obtain toe pressures ( $n=3$ ), leaving 415 patients/830 limbs eligible for further analysis. Of these patients, 210 (51%) had a resting  $TBI \leq 0.70$ , and 205 (49%) had a resting  $TBI > 0.70$ . Patient demographics for the two groups are presented in Table I. The parameters derived from pretest ankle and toe pressure assessment are presented in Table II, along with postexercise measurements. Overall, the patients with low TBIs had lower pre- and post-exercise mean ankle pressures, and mean ABI than patients with a normal TBI (all  $P \leq .002$ ).

#### 3.2 Diagnostic Agreement

A total of 117 (28%) of the 415 patients had PAD according to the outcome of the exercise testing and the TBI and 128 (31%) had normal test results for both evaluations according to AHA/ACC criteria. Another 93 (22%) patients had a low TBI but a normal treadmill testing result, and 77 (19%) patients had a normal TBI but PAD according to the treadmill testing. The overall agreement in the diagnostic classification was 245/415 (59%) with a Cohen's kappa ( $\kappa$ ) = 0.181 (95% CI: 0.087-0.276). When analysing the agreement on a limb basis, 160 (49%) of the 325 limbs with low TBI had an abnormal exercise test, whereas 165 (33%) of the 505 limbs with a normal TBI had an abnormal exercise test. The overall agreement on a limb basis was 500/830 (60%) with a Cohen's kappa ( $\kappa$ ) = 0.166 (95% CI: 0.096-0.232). The diagnostic outcome for the exercise testing on a limb basis is displayed in Table III and on a patient basis in Table IV. The sensitivity, specificity, positive and negative predictive values for the TBI with postexercise ABI as reference are shown in Table V. There were no significant differences in the ratio of abnormal tests within the two major

groups when comparing patients who were able to complete the full 5 minutes of the treadmill testing to the subgroups that were only able to complete less than 3 minutes or 3 to 5 minutes, respectively (all  $P \geq .481$ ). There were no significant differences in the ratio of abnormal exercise test results in the subgroup of patients with diabetes (49% abnormal tests) or chronic kidney failure (54% abnormal tests) compared to the other groups ( $P \geq .144$  for both).

### 3.3 The ABI and TBI versus outcome of exercise test

The probability of an abnormal treadmill testing result increased with the magnitude of the TBI reduction, as shown in Fig. 1 ( $P < .005$ ). The same was true for pre-exercise ABI (Fig. 2), with a borderline reduced ABI (0.91-1.00) having a higher probability (52%) of an abnormal exercise test than limbs with an ABI  $> 1.00$  (36%) ( $P < .0001$ ). The groups with a low TBI had a significantly higher ratio of abnormal test results than patients with a normal TBI, in limbs with ABI (0.96-1.00) and ABI  $> 1.10$  ( $P \leq .022$  for both). However, there were no statistically significant differences in the remaining groups ( $P \geq .200$  for all). The correlation between the pretest ABI and the TBI is shown in Fig. 3. A receiver operator curve for ABI and TBI was constructed and showed an area under the curve of 0.564 (95% CI: 0.509-0.619) for the ABI and 0.580 for the TBI (95% CI: 0.525-0.634), and no clear diagnostic cut-off was determined for the prediction of the outcome of exercise test (Fig. 4).

236

237 **4 Discussion**

238 This study shows that the TBI and postexercise ABI are not interchangeable methods for the  
239 diagnosis of PAD, with agreement in only 59% of cases. Although we found a correlation between  
240 the magnitude of the TBI decrease and the probability of an abnormal exercise test result, this could  
241 be partially explained by the known correlation between the resting ABI and TBI within the  
242 investigated pressure range.<sup>9</sup> The lower the resting ABI, the higher the probability of obtaining an  
243 abnormal test result, as was also indicated by this study.<sup>10</sup> Since limbs with a low TBI have a lower  
244 ABI in average than limbs with a normal TBI, there is a logical increase in the probability of an  
245 abnormal test result in that group. Accordingly, we did not detect a significant difference between  
246 patients with a normal TBI and low TBI within most of the ABI subgroups.

247

248 Having a reduced resting ABI is a well established indicator of increased risk of cardiovascular  
249 morbidity and mortality.<sup>3</sup> Furthermore, studies have shown that an abnormal postexercise ABI in  
250 the presence of a normal resting ABI, is an independent predictor of mortality and is related to a  
251 higher incidence of revascularization.<sup>11,12</sup> The TBI has also been shown to be an independent risk  
252 marker for mortality, although this remains to be verified in large-scale trials.<sup>5, 13,14</sup> It has been  
253 hypothesized that a reduced TBI in the presence of a normal ABI reflects small vessel disease, and  
254 the TBI has been shown to have a superior correlation to prognostic markers in comorbidities such  
255 as diabetes, kidney disease and microvascular disease than the ABI.<sup>5</sup> However, the group of patients  
256 with a low TBI and normal ABI likely contains a mixture of patients with masked large vessel  
257 disease (e.g., due to vessel stiffness), patients with small vessel disease, or the use of flawed  
258 diagnostic limits for the TBI. The treadmill testing could in theory offer a way to discriminate these

subgroups, as significant large vessel stenosis would result in a postexercise ABI decrease.<sup>15</sup>

Patients with diabetes are more prone to develop distal lesions, whereas smokers or young patients are more prone to develop proximal lesions.<sup>16</sup> However, we did not find any significant difference in the probability of an abnormal test result for patients with suspected microvascular disease, such as patients with diabetes or chronic kidney failure, although these subgroups were too small to allow any firm conclusions to be made.

Large-scale studies have shown, that a supranormal ABI is associated with increased mortality due to vessel stiffness.<sup>17</sup> Recently, Hammand and coworkers found that patients with an abnormal increase in ABI following an exercise test also have an increased risk of mortality, although this finding needs to be verified.<sup>8</sup> It could be hypothesized that this is a reflection of increased vessel stiffness which compromises the arterial windkessel function, leading to an alteration of the pressure curve.<sup>18</sup> However, we did not find any discrepancies in the ratio of patients with an abnormally high postexercise ABI between patients with a low or normal TBI.

In the 2016 ACC/AHA guidelines on the management of patients with lower extremity PAD, the postexercise ABI is recommended for patients with exertional nonjoint related symptoms and resting ABI within the range of 0.91-1.40 for the PAD diagnosis.<sup>2</sup> On the other hand, it is also stated that PAD can be diagnosed by measuring a TBI  $\leq 0.70$ , and this method is recommended in patients with a supranormal ABI ( $>1.40$ ) or patients with nonhealing wounds or gangrene. In other words, the methods are used more or less interchangeably to establish a PAD diagnosis. The findings in our study highlight that there is substantial disagreement between the two diagnostic modalities, likely due to differences in the sites of the vessel lesions. The treadmill testing primarily offers information on lesions proximal to the ankle, whereas patients with a reduced TBI are a

heterogeneous group including both patients with masked large vessel disease and patients with distal vessel lesions.<sup>16</sup> Another possible reason for the discrepancy is the limited evidence for the diagnostic limits in use for both methods.<sup>5,19</sup>. There is a substantial need for additional large-scale trials that correlate these methods to angiographically verified vessel stenoses and cardiovascular morbidity and mortality to clarify this.

## 5 Conclusion

The results of the present study show that the magnitude of the TBI reduction correlates with an increased probability of an abnormal postexercise ABI with 49% of limbs with a low TBI having an abnormal test result vs. 33% of the limbs with normal TBI. However, this was due in part to limbs with a low TBI having a lower resting ABI on average than limbs with a normal TBI, as the level of the resting ABI also relates to the probability of an abnormal exercise test result. This study shows that the TBI and the postexercise ABI are not interchangeable for establishing a PAD diagnosis, and likely reveal different entities of peripheral arteriosclerotic disease.

## 6 Acknowledgements

We would like to thank the laboratory technicians for their contributions to this study. This study received no financial support.

## 7 Conflict of Interest Statement

The authors report no conflicts of interest.

305

306

Journal Pre-proof

## 307 8 References

- 308 1. Norgren L, Hiatt WR, Dormandy JA, et al. Inter-Society Consensus for the Management of Peripheral  
 309 Arterial Disease (TASC II). *Eur J Vasc Endovasc Surg.* 2007;33 Suppl 1:S1-75.
- 310 2. Gerhard-Herman MD, Gornik HL, Barrett C, et al. 2016 AHA/ACC Guideline on the Management of  
 311 Patients With Lower Extremity Peripheral Artery Disease: Executive Summary: A Report of the American  
 312 College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation.*  
 313 2017;135:e686-e725.
- 314 3. Aboyans V, Criqui MH, Abraham P, et al. Measurement and interpretation of the ankle-brachial index: a  
 315 scientific statement from the American Heart Association. *Circulation.* 2012;126:2890-2909.
- 316 4. Suominen V, Rantanen T, Venermo M et al. J. Prevalence and risk factors of PAD among patients with  
 317 elevated ABI. *Eur J Vasc Endovasc Surg.* 2008;35:709-714.
- 318 5. Høyer C, Sandermann J, Petersen LJ. The toe-brachial index in the diagnosis of peripheral arterial disease.  
 319 *J Vasc Surg.* 2013;58:231-238.
- 320 6. Høyer C, Høgh AL, Sandermann J et al. Risk factors and haemodynamic variables in patients with low  
 321 toe-brachial index but normal ankle-brachial index. *Atherosclerosis* 2019;289:21-26.
- 322 7. Høyer C, Nielsen NS, Jordansen MKO et al. Comparison of two methods based on photoplethysmography  
 323 for the diagnosis of peripheral arterial disease. *Scand J Clin Lab Invest.* 2017;77:622-627.
- 324 8. Hammad TA, Hiatt WR, Gornik HL, et al. Prognostic value of an increase in post-exercise ankle-brachial  
 325 index. *Vasc Med.* 2017;22:204-209.

- 326 9. Aboyans V, Ho E, Denenberg JO et al. The association between elevated ankle systolic pressures and  
 327 peripheral occlusive arterial disease in diabetic and nondiabetic subjects. *J Vasc Surg.* 2008;48:1197-1203.
- 328 10. Armstrong DWJ, Tobin C, Matangi MF. Predictors of an abnormal postexercise ankle brachial index:  
 329 Importance of the lowest ankle pressure in calculating the resting ankle brachial index. *Clin Cardiol.*  
 330 2017;40:1163-1168.
- 331 11. Sheikh MA, Bhatt DL, Li J, et al. Usefulness of postexercise ankle-brachial index to predict all-cause  
 332 mortality. *Am J Cardiol.* 2011;107:778-782.
- 333 12. Hammad TA, Strefling JA, Zellers PR, et al. The Effect of Post-Exercise Ankle-Brachial Index on Lower  
 334 Extremity Revascularization. *JACC Cardiovasc Interv.* 2015;8:1238-1244.
- 335 13. Wickstrom JE, Laivuori M, Aro E, et al. Toe Pressure and Toe Brachial Index are Predictive of  
 336 Cardiovascular Mortality, Overall Mortality, and Amputation Free Survival in Patients with Peripheral  
 337 Artery Disease. *Eur J Vasc Endovasc Surg.* 2017;53:696-703.
- 338 14. Zobel EH, von Scholten BJ, Reinhard H, et al. Toe-brachial index as a predictor of cardiovascular  
 339 disease and all-cause mortality in people with type 2 diabetes and microalbuminuria. *Diabetologia.*  
 340 2017;60:1883-1891.
- 341 15. Tehan PE, Barwick AL, Sebastian M et al. Diagnostic accuracy of the postexercise ankle-brachial index  
 342 for detecting peripheral artery disease in suspected claudicants with and without diabetes. *Vasc Med.*  
 343 2018;23:116-125.
- 344 16. Aboyans V, Lacroix P, Criqui MH. Large and small vessels atherosclerosis: similarities and differences.  
 345 *Prog Cardiovasc Dis.* 2007;50:112-125.



- 346 17. Ankle Brachial Index Collaboration, Fowkes FG, Murray GD, et al. Ankle brachial index combined with  
347 Framingham Risk Score to predict cardiovascular events and mortality: a meta-analysis. *JAMA*.  
348 2008;300:197-208.
- 349 18. Westerhof N, Lankhaar JW, Westerhof BE. The arterial Windkessel. *Med Biol Eng Comput*.  
350 2009;47:131-141.
- 351 19. Mahe G, Pollak AW, Liedl DA, et al. Discordant Diagnosis of Lower Extremity Peripheral Artery  
352 Disease Using American Heart Association Postexercise Guidelines. *Medicine (Baltimore)*. 2015;94:e1277.
- 353

**Table I:** Demographics.

	<b>Normal TBI (n=205)</b>	<b>Low TBI (n=210)</b>	<b>P-value</b>
Age [years]	61.0 ± 12.4	65.9 ± 11.4	< .001
Height [cm]	173.5 ± 9.1	172.9 ± 9.7	.526
Weight [kg]	86.4 ± 18.5	82.6 ± 19.1	.043
Diabetes Mellitus	26 (13%)	32 (15%)	.453
Myocardial Infarction	21 (10%)	31 (15%)	.165
Chronic Kidney Insufficiency	7 (3%)	10 (5%)	.489
Arterial Hypertension	106 (52%)	107 (51%)	.878
Hypercholesterolemia	97 (47%)	107 (51%)	.459
Smokers	56 (27%)	48 (23%)	.295

*Abbreviations:* Data shown as mean (± standard deviation) or total (percentage).

**Table II:** Hemodynamic variables.

	Normal TBI (n=205)	Low TBI (n=210)	P-value
<b>Pre-test resting values</b>			
Brachial pressure [mmHg]	135 ± 16	136 ± 19	.534
Right ankle pressure [mmHg]	152 ± 21	146 ± 24	.002
Left ankle pressure [mmHg]	152 ± 22	146 ± 22	.005
Right ABI	1.13 ± 0.10	1.07 ± 0.09	< .001
Left ABI	1.12 ± 0.09	1.07 ± 0.09	< .001
Right toe pressure [mmHg]	113 ± 18	87 ± 21	< .001
Left toe pressure [mmHg]	114 ± 18	87 ± 20	< .001
Right TBI	0.84 ± 0.09	0.63 ± 0.12	< .001
Left TBI	0.84 ± 0.09	0.64 ± 0.12	< .001
<b>1-minute post-exercise</b>			
Brachial pressure [mmHg]	172 ± 26	176 ± 29	.182
Right ankle pressure [mmHg]	167 ± 39	150 ± 51	< .001
Left ankle pressure [mmHg]	167 ± 40	149 ± 48	< .001
Right ABI	0.98 ± 0.19	0.85 ± 0.24	< .001
Left ABI	0.97 ± 0.19	0.85 ± 0.23	< .001

*Abbreviations:* Data shown as mean (± standard deviation) or total (percentage).

**Table III:** Exercise testing on a limb basis.

	<b>Normal TBI (n=505 limbs)</b>	<b>Low TBI (n=325 limbs)</b>	<b>P-value</b>
≥30 mmHg decrease in ankle pressure	43 (9%)	59 (18%)	.0003
≥20% decrease in ABI	165 (33%)	160 (49%)	.002
≥30 mmHg decrease in ankle pressure or ≥20% decrease in ABI	165 (33%)	160 (49%)	.002
Postexercise increase in ABI	80 (16%)	47 (14%)	.644
Postexercise ABI drop 0-19%	260 (51%)	119 (37%)	.009

*Abbreviations:* Data shown as the mean ( $\pm$  standard deviation) or total (percentage).


**Table IV:** Exercise testing on a patient basis.

	<b>Normal TBI (n=205)</b>	<b>Low TBI (n=210)</b>	<b>P-value</b>
≥30 mmHg decrease in ankle pressure	24 (12%)	48 (23%)	.003
≥20% decrease in ABI	77 (38%)	117 (56%)	.0002
≥30 mmHg decrease in ankle pressure or ≥20% decrease in ABI	77 (38%)	117 (56%)	.0002
Postexercise increase in ABI	43 (21%)	39 (19%)	.539
Postexercise ABI drop 0-19%	87 (42%)	56 (27%)	.001


*Abbreviations:* Data shown as the mean ( $\pm$  standard deviation) or total (percentage).

**Table V:** Test accuracy of the TBI with postexercise ABI as reference.


	<b>On a patient basis (n=415)</b>	<b>On a limb basis (n=830)</b>
Sensitivity	60.3%	49.2%
Specificity	57.9%	67.3%
Positive predictive value	55.7%	49.2%
Negative predictive value	62.4%	67.3%




Downloaded for Anonymous User (n/a) at Aalborg Hospital from ClinicalKey.com by Elsevier on October 28, 2019.  
For personal use only. No other uses without permission. Copyright ©2019. Elsevier Inc. All rights reserved.



Downloaded for Anonymous User (n/a) at Aalborg Hospital from ClinicalKey.com by Elsevier on October 28, 2019.  
For personal use only. No other uses without permission. Copyright ©2019. Elsevier Inc. All rights reserved.







**Figure 1:** The probability of an abnormal exercise test vs. the TBI on a limb basis.

**Figure 2:** The probability of an abnormal exercise test vs. the ABI on a limb basis. \*) Denotes  $P \leq .022$

**Figure 3:** The correlation between TBI and pre-test ABI. The equation for the linear regression line (dotted line) were  $y = 0.63x + 0.01$  ( $R^2 = 0.173$ ). Full dots indicate patients with an abnormal exercise test and hollow dots patients with a normal exercise test.

**Figure 4:** Receiver operator characteristics (ROC) curve for various TBI (full line) and pretest ABI (dotted line) cut-offs.